

## **GCE AS/A Level**

0978/01



# MATHEMATICS – FP2 Further Pure Mathematics

MONDAY, 26 JUNE 2017 – AFTERNOON 1 hour 30 minutes

### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a WJEC pink 16-page answer booklet;
- · a Formula Booklet;
- a calculator.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer all questions.

Sufficient working must be shown to demonstrate the mathematical method employed.

## **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

CJ\*(S17-0978-01)

**1.** The function f is defined on the domain  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  by

$$f(x) = \sec x + x \tan x.$$

Determine whether f is even, odd or neither even nor odd.

[3]

[5]

2. Evaluate the integral

$$\int_0^2 \frac{2x^2 + 5}{x^2 + 4} \, \mathrm{d}x \, dx$$

giving your answer in the form  $a + b\pi$ , where a, b are constants to be determined.

- 3. Find the three cube roots of the complex number -8i. Give your answers in the form x + iy where x, y are either integers or surds. [8]
- 4. (a) Given that  $z = \cos \theta + i \sin \theta$ , show that

$$z^n + \frac{1}{z^n} = 2\cos n\theta$$

and find a similar expression for  $z^n - \frac{1}{z^n}$ .

[4]

[5]

(b) By expanding  $\left(z + \frac{1}{z}\right)^5$ , show that

$$\cos^5\theta = a\cos 5\theta + b\cos 3\theta + c\cos \theta,$$

where *a*, *b*, *c* are constants whose values should be determined.

(c) Hence evaluate the integral

$$\int_{0}^{\frac{\pi}{2}} \cos^{5}\theta \, \mathrm{d}\theta \, \cdot \tag{4}$$

5. Find the general solution to the equation

$$\cos\theta - \cos \theta = \sin 3\theta. \tag{8}$$

**6.** The function *f* is defined by

$$f(x) = \frac{24x^2 + 31x + 9}{(x+1)(2x+1)(3x+1)}$$

(a) Express f(x) in partial fractions.

[4]

(b) (i) Evaluate the integral

$$\int_0^2 f(x) dx$$

giving your answer as  $\ln N$ , where N is a positive integer.

(ii) Explain briefly why the integral

$$\int_{-2}^{0} f(x) \mathrm{d}x$$

cannot be evaluated.

[5]

- 7. (a) The point P(x, y) moves in such a way that its distance from the point (a, 0) is equal to its distance from the line x = -a. Show that the locus of P is the parabola with equation  $y^2 = 4ax$ .
  - (b) Determine the equation of the normal at the point  $(at^2, 2at)$  on the parabola. [4]
  - (c) This normal intersects the parabola again at the point  $(as^2, 2as)$ . Obtain an expression for s in terms of t. [5]
- **8.** The function f is defined by

$$f(x) = x + 3 + \frac{1}{x+1}$$
.

- (a) Find the equation of
  - (i) the vertical asymptote on the graph of f,
  - (ii) the asymptote that is not parallel to a coordinate axis.

[2]

- (b) Find the coordinates of the stationary points on the graph of f.
- [5]

[3]

- (c) (i) Obtain an expression for f''(x).
  - (ii) Hence classify each of the stationary points as a maximum or a minimum.
- (d) Sketch the graph of f, including the asymptotes. [3]
- (e) The set S is given by S = [4, 5]. Determine  $f^{-1}(S)$ . [4]